



## Table of Contents

### VOLUME 108

#### *Instructions to Authors*

v

#### ORIGINAL RESEARCH ARTICLES

Risk and safety of repetitive transcranial magnetic stimulation: report and suggested guidelines from the International Workshop on the Safety of Repetitive Transcranial Magnetic Stimulation, June 5–7, 1996 E.M. Wassermann (USA)	1
Usefulness of neurogenic motor evoked potentials for spinal cord monitoring findings in 112 consecutive patients undergoing surgery for spinal deformity Y. Péréon, J.-M. Bernard, G. Fayet, J. Delécrin, N. Passuti, P. Guihéneuc (France)	17
Motor and somatosensory evoked potentials in cervical spondylotic myelopathy A.M. de Noordhout, S. Myressiotis, V. Delvaux, J.D. Born, P.J. Delwaide (Belgium)	24
Multi-start downhill simplex method for spatio-temporal source localization in magnetoencephalography M. Huang, C.J. Aine, S. Supek, E. Best, D. Ranken, E.R. Flynn (USA, Croatia)	32
A MEG analysis of the P300 in visual discrimination tasks A. Mecklinger, B. Maess, B. Opitz, E. Pfeifer, D. Cheyne, H. Weinberg (Germany, Canada)	45
Somatosensory evoked magnetic fields elicited by dorsal penile, posterior tibial and median nerve stimulation H. Nakagawa, T. Namima, M. Aizawa, K. Uchi, Y. Kaiho, K. Yoshikawa, S. Orikasa, N. Nakasato (Japan)	57
Component wave analysis of flash visual evoked potentials in preterm infants K. Sawaguchi, T. Ogawa (Japan)	62
Impaired VEP after photostress response in multiple sclerosis patients previously affected by optic neuritis V. Parisi, F. Pierelli, R. Restuccia, M. Spadaro, L. Parisi, G. Colacino, M.G. Bucci (Italy)	73
Age and stage dependency of P300 latency alterations in non demented Parkinson's disease patients without therapy P. Stanzione, R. Semprini, M. Pierantozzi, A.M. Santilli, L. Fadda, R. Traversa, A. Peppe, G. Bernardi (Italy)	80
Generation of far field potentials from the trigeminal nerve in the cat Y. Ajimi, T. Ohira, T. Kawase, M. Takase (Japan)	92
Spontaneous EEG theta activity controls frontal visual evoked potential amplitudes E. Başar, E. Rahn, T. Demiralp, M. Schürmann (Germany, Turkey)	101
Localization of spatial attention processes with the aid of a probe technique K.J. Bruin, J.L. Kenemans, M.N. Verbaten, A.H.C. Van der Heijden (The Netherlands)	110
A method for the calculation of induced band power: implications for the significance of brain oscillations W. Klimesch, H. Russegger, M. Doppelmayr, Th. Pachinger (Austria)	123
Auditory evoked potentials to abrupt pitch and timbre change of complex tones: electrophysiological evidence of 'streaming'? S.J. Jones, O. Longe, M. Vaz Pato (UK)	131
Impaired mismatch negativity (MMN) generation in schizophrenia as a function of stimulus deviance, probability, and interstimulus/interdeviant interval D.C. Javitt, S. Grochowski, A.-M. Shelley, W. Ritter (USA)	143
The mismatch negativity for duration decrement of auditory stimuli in healthy subjects S.-L. Joutsiniemi, T. Ilvonen, J. Sinkkonen, M. Huottilainen, M. Tervaniemi, A. Lehtokoski, T. Rinne, R. Näätänen (Finland)	154
Differential effects of normal aging on sources of standard N1, target N1 and target P300 auditory event-related brain potentials revealed by low resolution electromagnetic tomography (LORETA) P. Anderer, R.D. Pascual-Marqui, H.V. Semlitsch, B. Saletu (Austria, Switzerland)	160
Isoflurane plus nitrous oxide versus propofol for recording of motor evoked potentials after high frequency repetitive electrical stimulation U. Pechstein, J. Nadstawek, J. Zentner, J. Schramm (Germany)	175
High-frequency (600 Hz) SEP activities originating in the subcortical and cortical human somatosensory system R. Gobbelé, H. Buchner, G. Curio (Germany)	182

Dissociation induced by voluntary movement between two different components of the centro-parietal P40 SEP to tibial nerve stimulation M. Valeriani, D. Restuccia, V. Di Lazzaro, C. Barba, D. Le Pera, P. Tonali (Italy)	190
Multimodality evoked potentials as a prognostic tool in term asphyxiated newborns E. Scalais, A. François-Adant, C. Nuttin, A. Bachy, J.M. Guerit (Belgium)	199
Chemosensory event-related potentials change with age T. Hummel, S. Barz, E. Pauli, G. Kobal (Germany)	208
Hemispheric asymmetry of the auditory evoked N100m response in relation to the crossing point between the central sulcus and Sylvian fissure S. Ohtomo, N. Nakasato, A. Kanno, K. Hatanaka, R. Shirane, K. Mizoi, T. Yoshimoto (Japan)	219
Presurgical functional localization of primary somatosensory cortex by dipole tracing method of scalp-skull-brain head model applied to somatosensory evoked potential S. Mine, N. Oka, A. Yamaura, Y. Nakajima (Japan)	226
Cortical localisation of magnetic fields evoked by oesophageal distension P.L. Furlong, Q. Aziz, K.D. Singh, D.G. Thompson, A. Hobson, G.F.A. Harding (UK)	234
Quantitative trait loci analysis of human event-related brain potentials: P3 voltage H. Begleiter, B. Porjesz, T. Reich, H.J. Edenberg, A. Goate, J. Blangero, L. Almasy, T. Foroud, P. Van Eerdewegh, J. Polich, J. Rohrbach, S. Kuperman, L.O. Bauer, S.J. O'Connor, D.B. Chorlian, T.K. Li, P.M. Conneally, V. Hesselbrock, J.P. Rice, M.A. Schuckit, R. Cloninger, J. Nurnberger, Jr., R. Crowe, F.E. Bloom (USA)	244
Dynamic spectral analysis of event-related potentials D. Melkonian, E. Gordon, C. Rennie, H. Bahramali (Australia)	251
P3, Positive slow wave and working memory load: a study on the functional correlates of slow wave activity L. García-Larrea, G. Cézanne-Bert (France)	260
Language-related hemispheric asymmetry in healthy subjects and patients with temporal lobe epilepsy as studied by event-related brain potentials and intracarotid amobarbital test W. Gerschlag, W. Laluschek, J. Lehrner, C. Baumgartner, G. Lindinger, W. Lang (Austria)	274
Auditory event-related potentials to semantic priming during sleep J. Brualla, M.F. Romero, M. Serrano, J.R. Valdizán (Spain)	283
Replicability of MEG and EEG measures of the auditory N1/N1m-response J. Virtanen, J. Ahveninen, R.J. Ilmoniemi, R. Näätänen, E. Pekkonen (Finland)	291
Topography and morphology of heart action-related EEG potentials G. Dirlich, T. Dietl, L. Vogl, F. Strian (Germany)	299
Cortical and thalamic visual evoked potentials during sleep-wake states and spike-wave discharges in the rat H.K.M. Meeren, E.L.J.M. Van Luijcklaar, A.M.L. Coenen (The Netherlands)	306
Single-sweep cortical somatosensory evoked potentials: N20 and evoked bursts in sevoflurane anaesthesia V. Jäntti, E. Sonkajärvi, S. Mustola, S. Rytty, P. Kiiski, K. Suominen (Finland)	320
Waveform analysis of spinal somatosensory evoked potential: paradoxically enhanced negative peaks immediately caudal to the site of conduction block T. Tani, T. Ushida, H. Yamamoto, J. Kimura (Japan)	325
Refractory properties of auditory brain-stem responses evoked by electrical stimulation of human cochlear nucleus: evidence of neural generators M.D. Waring (USA)	331
Interstimulus interval and auditory event-related potentials in children: evidence for multiple generators R. Čeponienė, M. Cheour, R. Näätänen (Finland)	345
The AEP T-complex to synthesised musical tones: left-right asymmetry in relation to handedness and hemisphere dominance S.J. Jones, C. Byrne (UK)	355
Automatic and effortful processes in auditory memory reflected by event-related potentials. Age-related findings E. Amenedo, F. Díaz (Spain)	361
Combined mapping of human auditory EEG and MEG responses M. Huotilainen, I. Winkler, K. Alho, C. Escera, J. Virtanen, R.J. Ilmoniemi, I.P. Jääskeläinen, E. Pekkonen, R. Näätänen (Finland, Hungary, Spain)	370
ERP components in category matching tasks J. Jun, B. Porjesz, H. Begleiter (USA, China)	380
Enhanced negative slope of the readiness potential preceding a target force production task H. Masaki, N. Takasawa, K. Yamazaki (Japan)	390
Very late pain-related activity identified with topographically mapped frequency domain analysis of evoked potentials D.E. Becker, R.S. Noss, G. Fein, C.D. Yingling (USA, Switzerland)	398
Three-dimensional tomography of event-related potentials during response inhibition: evidence for phasic frontal lobe activation W.K. Strik, A.J. Fallgatter, D. Brandeis, R.D. Pascual-Marqui (Germany, Switzerland)	406



The scalp to earlobe montage as standard in routine SEP recording. Comparison with the non-cephalic reference in patients with lesions of the upper cervical cord M. Valeriani, D. Restuccia, V. Di Lazzaro, D. Le Pera, C. Barba, P. Tonali (Italy)	414
Dermatomal and mixed nerve somatosensory evoked potentials in the diagnosis of neurogenic thoracic outlet syndrome R. Cakmur, F. Idiman, E. Akalin, A. Genç, G.G. Yener, V. Öztürk (Turkey)	423
The covert interplay between perception and memory: event-related potential evidence M.P. Viggiano, M. Kutas (Italy, USA)	435
The N18 component of the median nerve SEP is not reduced by vibration G.M. Manzano, N. Negrão, J.A.M. Nóbrega (Brazil)	440
Magnetic stimulation of the corticospinal tracts in Pelizaeus-Merzbacher disease A. Nezu, S. Kimura, S. Takeshita, H. Osaka, M. Tanaka (Japan)	446
Relation of olfactory event-related potentials to changes in stimulus concentration T. Tateyama, T. Hummel, S. Roscher, H. Post, G. Kobal (Japan, USA, Germany)	449
The N400 in a semantic categorization task across 6 decades M. Kutas, V. Iragui (USA)	456
Visual cognitive dysfunction in depression: an event-related potential study F. Bange, N. Bathien (France)	472
Sympathetic skin responses recorded from non-palmar and non-plantar skin sites: their role in the evaluation of thermal sweating K. Matsunaga, T. Uozumi, S. Tsuji, Y. Murai (Japan)	482
Neural conduction in visual pathways in newly-diagnosed IDDM patients V. Parisi, L. Uccioli, L. Parisi, G. Colacino, G. Manni, G. Menzinger, M.G. Bucci (Italy)	490
EEG theta and frontal alpha oscillations during auditory processing change with aging J.Y. Yordanova, V.N. Kolev, E. Başar (Bulgaria, Germany)	497
Spike-induced interference in auditory sensory processing in Landau-Kleffner syndrome S. Seri, A. Cerquiglioni, F. Pisani (Italy)	506
Effects of localized pontine lesions on auditory brain-stem evoked potentials and binaural processing in humans H. Pratt, A. Polyakov, V. Aharonson, A.D. Korczyn, R. Tadmor, B.C. Fullerton, R.A. Levine, M. Furst (Israel)	511
Cochlear implant performance and electrically-evoked auditory brain-stem response characteristics S. Gallégo, B. Frachet, C. Micheyl, E. Truy, L. Collet (France)	521
Quantification and rejection of ocular artifacts in auditory evoked fields in schizophrenics T. Meier, T. Rosburg, M. Arnold, I. Kreitschmann-Andermahr, H. Sauer, H. Nowak, H. Witte (Germany)	526
High frequency oscillations in early cortical somatosensory evoked potentials I. Ozaki, C. Suzuki, Y. Yaegashi, M. Baba, M. Matsunaga, I. Hashimoto (Japan)	536
Evidence for efferent effects on early components of the human auditory brain-stem evoked potentials A. Polyakov, H. Pratt, Y. Shi (Israel, China)	543
P300 and stress in mild head injury patients Y. Granovsky, E. Sprecher, J. Hemli, D. Yarnitsky (Israel)	554
Identification of the stimulated hemiretinae using a reduced number of PVEP trials K. Hoffmann, H. Witte, H. Niedner, R. Vollandt (Germany)	560
Event-related potentials in patients with temporal lobe epilepsy reveal topography specific lateralization in relation to the side of the epileptic focus W. Lalouschek, W. Gerschlagel, J. Lehrner, C. Baumgartner, G. Lindinger, L. Deecke, W. Lang (Austria)	567
Comparison of distortion product otoacoustic emissions with auditory brain-stem response for clinical use in neonatal intensive care unit A. Ochi, A. Yasuhara, Y. Kobayashi (Japan)	577
P9 in median nerve SEPs is a junctional potential generated by the change of the volume conductor size between trunk and neck A. Mochizuki, M. Sonoo, T. Shimizu, M. Iwata (Japan)	584
Brain electrical source analysis of primary cortical components of the tibial nerve somatosensory evoked potential using regional sources U. Baumgärtner, H. Vogel, J. Ellrich, J. Gawehn, P. Stoeter, R.-D. Treede (Germany)	588
<i>Index of Authors to Volume 108</i>	600
<i>Index of Subjects to Volume 108</i>	605
<i>Table of Contents to Volume 108</i>	613



